

FINAL EXAM
Please do individually

1. Long gravity waves obey the dispersion relation

$$\omega^2 = gH(k^2 + l^2)$$

Where H is the water depth and g is the gravity acceleration. Suppose that a packet of waves with $\mathbf{k}=(k_0, l_0)$ propagates from a region of uniform depth H_0 into water which slowly deepens as x increases, but the depth is independent of y .

- a) Find the depth to which the waves can propagate. (i.e., where is the turning point where $c_{gx}=0$)?
 - b) Find the shape of the ray (function of (x, y)) in the vicinity of the turning depth H_T . (hint: Taylor expand in the neighborhood of H_T .)
2. Now consider the same packet from problem 1 to propagate into water that slowly shallows as x increases. The waves have an initial amplitude of η_0 in the region prior to the sloping topography (where $H=H_0$).
- a) Find the relation for the variation of wave amplitude as the waves move into shallower water. When the water depth becomes very small how does the amplitude depend on H ? (you will need the energy equation).
 - b) How does the amplitude behave in the neighborhood of the turning point in problem 1?

You will need to use the fact that energy, averaged over a wave period, or wavelength, is $E=\rho g \eta^2/2$ and assume that the topography is so weakly varying that reflection is unimportant.

3. Consider a nonrotating stratified fluid with constant N . At time $t=0$ a small region of the fluid is disturbed near the origin $x=0, z=0$. This generates internal waves that radiate away from the disturbance site. The disturbance generates a broad band of wave frequencies, but the wavelengths are limited to a maximum of λ_m .
- a) If the fluid is taken to be inviscid find the region of the fluid that could contain waves at some later time $t > 0$. For simplicity take the flow to be two-dimensional (x, z) . Consider the trajectory of the ray emanating from the source at $(0,0)$.
 - b) After a long time the effects of viscosity can be expected to damp the waves. For constant viscosity ν the equation governing linear internal waves is (in (x, z)).


$$\nabla^2 w_{tt} + N^2 w_{xx} = u \nabla^4 w_t$$

From this equation determine the dispersion relation. Examine the limit where $[\nu(k^2+m^2)]^2 \gg 4N^2 \cos^2 \theta$. To which type of regime does this limit correspond? Here $\cos \theta = k / (k^2+m^2)^{1/2}$ and k and m are the horizontal and vertical wavenumbers, respectively. Write the resulting solution. When does the minimum decay occur? Now examine the opposite limit $[\nu(k^2+m^2)]^2 \ll 4N^2 \cos^2 \theta$ and write the corresponding solution. To which type of regime does this limit correspond?

- Consider a layer of incompressible fluid of mean depth D bounded above by a rigid lid, and rotating with a constant frequency $f/2$. For $y < 0$, $f \nabla h_B / D = \beta_1 \hat{j}$ and for $y > 0$, $f \nabla h_B / D = \beta_2 \hat{j}$, where β_1 and β_2 are constants. In the context of linear quasi-geostrophic theory discuss what happens to a wave with wavenumber $k_I = \hat{i} k_I + \hat{j} l_I$ incident upon the line $y=0$ from the region $y < 0$. Find the reflected and transmitted wave amplitudes and wavenumbers. Note that β_1 and β_2 are not equal. When is the reflection total? (Hint: what is preserved in the reflection/transmission?)
- Consider a layer of uniform density on the beta-plane ($f=f_0+\beta y$) and mean thickness H . The motion is governed by the following equation:

$$\begin{aligned} fu &= -g\eta_y \\ fv &= g\eta_x \\ \eta_t + H(u_x + v_y) &= -\lambda\eta + W(x, y, t). \end{aligned}$$

Where η is the departure of the layer depth from the rest value H , λ is a constant representing the effects of mixing, W is a prescribed vertical mass flux. At $t=0$ the vertical mass flux $W=w_0\delta(x-x_0)$ for $y_S < y < y_N$ and $W=0$ elsewhere is tuned on and held fixed, dividing the (x, y) plane in different regions as sketched below:



Find the resulting flow in each of the above regions. Interpret your results in terms of Rossby wave dynamics: which type of Rossby wave do you have? Consider a time $t_1 > 0$: how far has the wave propagated at y_S , y_N and in between?

6. Consider a two-layer dry compressible atmosphere. The lower layer, $0 < z < D$, is isothermal with temperature T_1 , and the upper layer, $D < z < \infty$ has temperature T_2 , also a constant. For each layer $N_1^2 = \text{constant}$; $N_2^2 = \text{constant}$.

- a) Find the basic state density and pressure profiles given the density $\rho = \rho_0$ at $z=0$. What is the density jump at $z=D$? What is the condition to have a stable atmosphere?
- b) Under what conditions could a linear baroclinic (i.e., vertical propagation) Rossby wave (with horizontal and temporal structure $\sim e^{i(kx+ly-\sigma t)}$) exist? You will need the vertical structure equation derived in class for both layers but do not need to solve the complete problem to give an answer.
- c) Now solve the problem for waves trapped in both layers ($q_1^2 < 0$; $q_2^2 < 0$). Specify the vertical structure from the vertical structure equation in both layers. You need to use the boundary conditions at $z=0$; $z=D$; $z \rightarrow +\infty$ (express the pressures and vertical velocities in terms of the stream function as done in class). Without fully solving the equations obtained from the boundary conditions, solve them for $\lambda=0$: what mode are you obtaining? What are the vertical profiles of $\varphi_1(z)$, $\varphi_2(z)$ in this case?

7. Consider a two-dimensional basin with homogeneous density ($\rho = \rho_0 = \text{constant}$, of depth D and length L in x -direction. The basin infinitely long in y , i.e. $\frac{\partial}{\partial y} = 0$

- a) Find the linear free surface gravity oscillation and the dispersion relationship. Describe the corresponding surface displacement patterns.
- b) Even though the general solution is a linear superposition of all the normal modes of the basin, consider one individual mode. Now a travelling pressure disturbance is forcing the basin

$$P_a = P_0 - \cos(kx - \omega t) = P_0 \operatorname{Re} \left[e^{i(kx - \omega t)} \right]$$

The following pressure fixes only the frequency response of the basin through the surface boundary condition on the velocity potential. Find the amplitude A_m of the n -normal mode. For what value of ω will P_a produce a layer response? How will the response depend on K ?

MIT OpenCourseWare
<http://ocw.mit.edu>

12.802 Wave Motion in the Ocean and the Atmosphere
Spring 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.