

8. LIMITS

Definition 8.1. Let $P \in \mathbb{R}^n$ be a point. The **open ball of radius** $\epsilon > 0$ **about** P is the set

$$B_\epsilon(P) = \{Q \in \mathbb{R}^n \mid \|\overrightarrow{PQ}\| < \epsilon\}.$$

The **closed ball of radius** $\epsilon > 0$ **about** P is the set

$$\{Q \in \mathbb{R}^n \mid \|\overrightarrow{PQ}\| \leq \epsilon\}.$$

Definition 8.2. A subset $A \subset \mathbb{R}^n$ is called **open** if for every $P \in A$ there is an $\epsilon > 0$ such that the open ball of radius ϵ about P is entirely contained in A ,

$$B_\epsilon(P) \subset A.$$

We say that B is **closed** if the complement of B is open.

Put differently, an open set is a union of open balls. Open balls are open and closed balls are closed. $[0, 1)$ is neither open nor closed.

Definition 8.3. Let $B \subset \mathbb{R}^n$. We say that $P \in \mathbb{R}^n$ is a **limit point** if for every $\epsilon > 0$ the intersection

$$B_\epsilon(P) \cap B \neq \emptyset.$$

Example 8.4. 0 is a limit point of

$$\left\{ \frac{1}{n} \mid n \in \mathbb{N} \right\} \subset \mathbb{R}.$$

Lemma 8.5. A subset $B \subset \mathbb{R}^n$ is closed if and only if B contains all of its limit points.

Example 8.6. $\mathbb{R}^n - \{0\}$ is open. One can see this directly from the definition or from the fact that the complement $\{0\}$ is closed.

Definition 8.7. Let $A \subset \mathbb{R}^n$ and let $P \in \mathbb{R}^n$ be a limit point. Suppose that $f: A \rightarrow \mathbb{R}^m$ is a function.

We say that f approaches L as Q approaches P and write

$$\lim_{Q \rightarrow P} f(Q) = L,$$

if for every $\epsilon > 0$ we may find $\delta > 0$ such that whenever $Q \in B_\delta(P) \cap A$ $f(Q) \in B_\epsilon(L)$. In this case we call L the **limit**.

It might help to understand the notion of a limit in terms of a game played between two people. Let's call the first player Larry and the second player Norman. Larry wants to show that L is the limit of $f(Q)$ as Q approaches P and Norman does not.

So Norman gets to choose $\epsilon > 0$. Once Norman has chosen $\epsilon > 0$, Larry has to choose $\delta > 0$. The smaller that Norman chooses $\epsilon > 0$,

the harder Larry has to work (typically he will have to make a choice of $\delta > 0$ very small).

Proposition 8.8. *Let $f: A \rightarrow \mathbb{R}^m$ and $g: A \rightarrow \mathbb{R}^m$ be two functions. Let $\lambda \in \mathbb{R}$ be a scalar. If P is a limit point of A and*

$$\lim_{Q \rightarrow P} f(Q) = L \quad \text{and} \quad \lim_{Q \rightarrow P} g(Q) = M,$$

then

- (1) $\lim_{Q \rightarrow P} (f + g)(Q) = L + M$, and
- (2) $\lim_{Q \rightarrow P} (\lambda f)(Q) = \lambda L$.

Now suppose that $m = 1$.

- (3) $\lim_{Q \rightarrow P} (fg)(Q) = LM$, and
- (4) if $M \neq 0$, then $\lim_{Q \rightarrow P} (f/g)(Q) = L/M$.

Proof. We just prove (1). Suppose that $\epsilon > 0$. As L and M are limits, we may find δ_1 and δ_2 such that, if $\|Q - P\| < \delta_1$ and $Q \in A$, then $\|f(Q) - L\| < \epsilon/2$ and if $\|Q - P\| < \delta_2$ and $Q \in A$, then $\|g(Q) - M\| < \epsilon/2$.

Let $\delta = \min(\delta_1, \delta_2)$. If $\|Q - P\| < \delta$ and $Q \in A$, then

$$\begin{aligned} \|(f + g)(Q) - L - M\| &= \|(f(Q) - L) + (g(Q) - M)\| \\ &\leq \|f(Q) - L\| + \|g(Q) - M\| \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon, \end{aligned}$$

where we applied the triangle inequality to get from the second line to the third line. This is (1). (2-4) have similar proofs. \square

Definition 8.9. *Let $A \subset \mathbb{R}^n$ and let $P \in A$. If $f: A \rightarrow \mathbb{R}^m$ is a function, then we say that f is continuous at P , if*

$$\lim_{Q \rightarrow P} f(Q) = f(P).$$

We say that f is continuous, if it is continuous at every point of A .

Theorem 8.10. *If $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is a polynomial function, then f is continuous.*

A similar result holds if f is a rational function (a quotient of two polynomials).

Example 8.11. *$f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $f(x, y) = x^2 + y^2$ is continuous.*

Sometimes Larry is very lucky:

Example 8.12. Does the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x - y},$$

exist? Here the domain of f is

$$A = \{(x, y) \in \mathbb{R}^2 \mid x \neq y\}.$$

Note $(0, 0)$ is a limit point of A . Note that if $(x, y) \in A$, then

$$\frac{x^2 - y^2}{x - y} = x + y,$$

so that

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x - y} = \lim_{(x,y) \rightarrow (0,0)} x + y = 0.$$

So the limit does exist.

Norman likes the following result:

Proposition 8.13. Let $A \subset \mathbb{R}^n$ and let $B \subset \mathbb{R}^m$. Let $f: A \rightarrow B$ and $g: B \rightarrow \mathbb{R}^l$.

Suppose that P is a limit point of A , L is a limit point of B and

$$\lim_{Q \rightarrow P} f(Q) = L \quad \text{and} \quad \lim_{M \rightarrow L} g(M) = E.$$

Then

$$\lim_{Q \rightarrow P} (g \circ f)(Q) = E.$$

Proof. Let $\epsilon > 0$. We may find $\delta > 0$ such that if $\|M - L\| < \delta$, and $M \in B$, then $\|g(M) - E\| < \epsilon$. Given $\delta > 0$ we may find $\eta > 0$ such that if $\|Q - P\| < \eta$ and $Q \in A$, then $\|f(Q) - L\| < \delta$. But then if $\|Q - P\| < \eta$ and $Q \in A$, then $M = f(Q) \in B$ and $\|M - L\| < \delta$, so that

$$\begin{aligned} \|(g \circ f)(Q) - E\| &= \|g(f(Q)) - E\| \\ &= \|g(M) - E\| \\ &< \epsilon. \end{aligned}$$

□

Example 8.14. Does

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

exist? The answer is no.

To show that the answer is no, we suppose that the limit exists. Suppose we consider restricting to the x -axis. Let

$$f: \mathbb{R} \rightarrow \mathbb{R}^2,$$

be given by $t \rightarrow (t, 0)$. As f is continuous, if we compose we must get a function with a limit,

$$\lim_{t \rightarrow 0} \frac{0}{t^2 + 0} = \lim_{t \rightarrow 0} 0 = 0.$$

Now suppose that we restrict to the line $y = x$. Now consider the function

$$f: \mathbb{R} \rightarrow \mathbb{R}^2,$$

be given by $t \rightarrow (t, t)$. As f is continuous, if we compose we must get a function with a limit,

$$\lim_{t \rightarrow 0} \frac{t^2}{t^2 + t^2} = \lim_{t \rightarrow 0} \frac{1}{2} = \frac{1}{2}.$$

The problem is that the limit along two different lines is different. So the original limit cannot exist.

Example 8.15. Does the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3}{x^2 + y^2},$$

exist? Let us use polar coordinates. Note that

$$\frac{x^3}{x^2 + y^2} = \frac{r^3 \cos^3 \theta}{r^2} = r \cos^3 \theta.$$

So we guess the limit is zero.

$$\begin{aligned} \lim_{(x,y) \rightarrow (0,0)} \left| \frac{x^3}{x^2 + y^2} \right| &= \lim_{r \rightarrow 0} |r \cos^3 \theta| \\ &\leq \lim_{r \rightarrow 0} |r| = 0. \end{aligned}$$

Example 8.16. Does the limit

$$\lim_{(x,y,z) \rightarrow (0,0,0)} \frac{xyz}{x^2 + y^2 + z^2},$$

exist? Same trick, but now let us use spherical coordinates.

$$\begin{aligned} \lim_{(x,y,z) \rightarrow (0,0,0)} \left| \frac{xyz}{x^2 + y^2 + z^2} \right| &= \lim_{\rho \rightarrow 0} \left| \frac{\rho^3 \cos^2 \phi \sin \phi \cos \theta \sin \theta}{\rho^2} \right| \\ &= \lim_{\rho \rightarrow 0} |\rho \cos^2 \phi \sin \phi \cos \theta \sin \theta| \\ &\leq \lim_{\rho \rightarrow 0} |\rho| = 0. \end{aligned}$$

Sometimes Norman needs to restrict to more complicated curves than just lines:

Example 8.17. *Does the limit*

$$\lim_{(x,y) \rightarrow (0,0)} \frac{y}{y + x^2},$$

exist? If we restrict to the line $t \rightarrow (at, bt)$, then we get

$$\lim_{t \rightarrow 0} \frac{bt}{bt + a^2t^2} = \lim_{t \rightarrow 0} \frac{b}{b + at} = 1.$$

But if we restrict to the conic $t \rightarrow (t, at^2)$, then we get

$$\lim_{t \rightarrow 0} \frac{at^2}{at^2 + t^2} = \lim_{t \rightarrow 0} \frac{a}{1 + a} = \frac{a}{1 + a},$$

and the limit changes as we vary a , so that the limit does not exist.

Note that if we start with

$$\frac{y}{y + x^d},$$

then Norman even needs to use curves of degree d ,

$$t \rightarrow (t, at^d).$$

MIT OpenCourseWare
<http://ocw.mit.edu>

18.022 Calculus of Several Variables
Fall 2010

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.