

18.311 — MIT (Spring 2014)

Rodolfo R. Rosales

March 16, 2014.

Problem Set # 04. Due: Friday April 4.

IMPORTANT:

- Turn in the regular and the special problems **stapled in two SEPARATE** packages.
- **PRINT your name** in each page of your answers. ***PRINT*** (the pencil will not break).
- In page one of each package **print the names** of the other members of your group.

Contents

1 Regular Problems.	1
1.1 Statement: Haberman problem 69.01	1
Rate of change of density for moving observer.	1
1.2 Statement: TFPa11. Longest queue through a light	2
Last car to make green light. Longest queue through a light.	2
1.3 DiAn21 statement: Non-dimensional form.	2
1.4 Statement: TFPb09. Quasi-linear equation solution	2
Solve a simple quasi-linear problem using characteristics.	2
1.5 Statement: TFPa21. Characteristics for 3D scalar quasi-linear equation	2
Write the characteristic equations for 3-D scalar quasi-linear equations.	2
Find the general solution for a simple linear equation.	2
1.6 Statement: TFPa18. Envelopes for families of curves	3
2 Special Problems.	3
2.1 DiAn26 statement: Non-dimensional form.	3
2.2 Statement: TFPa20. Semi-linear 1 st order eqn. & characteristics.	4

1 Regular Problems.

1.1 Statement: Haberman problem 69.01.

Show that for an *observer moving with the traffic*, the rate of change of the measured density is

$$\frac{d\rho}{dt} = (u - c(\rho)) \rho_x, \tag{1.1}$$

where $c = \frac{dq}{d\rho}$.

1.2 Statement: TFPa11. Longest queue through a light.

A traffic signal (at $x = 0$) is green for $0 \leq t \leq T$, and red for all other times. If $\rho(x, 0) = \rho_j$ for $x \leq 0$, $\rho(x, 0) = 0$ for $x > 0$, and $q = (4q_m/\rho_j^2)\rho(\rho_j - \rho)$, determine the trajectory of the last car to make the light. What is the longest traffic queue that can pass through the intersection during the green light?

1.3 DiAn21 statement: Non-dimensional form.

Consider the problem (5th order Linear KDV equation Initial Value Problem)

$$\frac{\partial \tilde{u}}{\partial \tilde{t}} + \alpha \frac{\partial \tilde{u}}{\partial \tilde{x}} + \beta \frac{\partial^5 \tilde{u}}{\partial \tilde{x}^5} = 0, \quad -\infty < x < \infty \text{ and } t > 0, \quad \tilde{u}(\tilde{x}, 0) = \frac{u_m d^2}{(d^2 + \tilde{x}^2)}, \quad (1.2)$$

where (a) $\alpha > 0$, $\beta > 0$, $u_m > 0$, and $d > 0$ are dimensional constants; (b) \tilde{u} is density of some quantity (wiggies per unit length); and (c) tildes (i.e.: \tilde{u} , \tilde{x} , and \tilde{t}) denote dimensional variables.

1. What are the dimensions of the constants α , β , u_m , and d ?

2. Introduce a-dimensional variables¹ u , x ,

and t , so that the equation takes the form $u_t + u_x + \gamma u_{xxxxx} = 0$, (1.3)

where γ is a constant without dimensions, and the initial condition involves no free constants.²

1.4 Statement: TFPb09. Quasi-linear equation solution.

Find the solution to

$$x^2 \psi_x - \psi^2 \psi_y = 0, \quad (1.4)$$

with $\psi = x$ on $y = x$, for $x > 0$. Where is the solution defined, and why?

1.5 Statement:

TFPa21. Characteristics for 3D scalar quasi-linear equation.

Consider a p.d.e. of the form

$$P(x, y, z, \psi) \psi_x + Q(x, y, z, \psi) \psi_y + R(x, y, z, \psi) \psi_z = W(x, y, z, \psi), \quad (1.5)$$

¹ That is, $x = \tilde{x}/L$, $t = \tilde{t}/T$, and $u = \tilde{u}/U$, for appropriate choices of a length L , a time T , and a density U .

² That is, no letter constants in it.

for the real valued function $\psi = \psi(x, y, z)$, for some given coefficient functions P, Q, R , and W . Assume that the values of ψ are given on some surface Γ . Specifically, let the surface be described (parametrically) by

$$x = X(u, v), \quad y = Y(u, v), \quad \text{and} \quad z = Z(u, v), \quad (1.6)$$

by some functions X, Y , and Z , defined in some region Γ_p of the u - v plane. Then

$$\psi = \Psi(u, v), \quad (1.7)$$

for some function Ψ .

(a) Give a detailed description of how you would solve the problem for ψ above, using the method of characteristics.

(b) Use your method to find the solution to

$$A\psi_x + B\psi_y + \psi_z = -\psi, \quad (1.8)$$

where A and B are constants, and $\psi = \Psi(x, y)$ for $z = 0$.

1.6 Statement: TFPa18. Envelopes for families of curves.

Find the envelope of each of the following family of curves: $\left\{ \begin{array}{l} \text{a. } y = cx - (1 + c^2)x^2. \\ \text{b. } x^2 + a^2y^2 = a. \\ \text{c. } (1 - c)x + cy = c - c^2. \end{array} \right.$

2 Special Problems.

2.1 DiAn26 statement: Non-dimensional form.

Consider the problem (Burgers' equation Boundary Value Problem)

$$\frac{\partial \tilde{u}}{\partial \tilde{t}} + \alpha \tilde{u} \frac{\partial \tilde{u}}{\partial \tilde{x}} - \beta \frac{\partial^2 \tilde{u}}{\partial \tilde{x}^2} = 0, \quad 0 < x < d \text{ and } t > 0, \quad \tilde{u}(0, \tilde{t}) = \tilde{u}(d, \tilde{t}) = 0, \quad (2.9)$$

with some initial data. Here (a) $\alpha > 0, \beta > 0$, and $d > 0$ are dimensional constants; (b) \tilde{u} is density of some quantity (wiggies per unit length); and (c) tildes (i.e.: \tilde{u}, \tilde{x} , and \tilde{t}) denote dimensional variables.

1. What are the dimensions of the constants α, β , and d ?

2. Introduce a-dimensional variables³ u, x , and t , so that the equation takes the form

$$u_t + u u_x - u_{xx} = 0, \quad (2.10)$$

where $0 < x < 1$.

³ That is, $x = \tilde{x}/L, t = \tilde{t}/T$, and $u = \tilde{u}/U$, for appropriate choices of a length L , a time T , and a density U .

2.2 Statement: TFPa20. Semi-linear 1st order eqn. & characteristics.

Consider the equation

$$x^2 \psi_x - x y \psi_y = \psi^2, \quad (2.11)$$

subject to $\psi = 1$ on the curve Γ given by $\mathbf{x} = \mathbf{y}^2$. This is a semi-linear problem (the terms involving derivatives of the solution are linear), that can be written in terms of characteristics.

- A.** Compute the characteristic curves that cross the curve Γ , as follows: (i) Parameterize the curve Γ , say: $\mathbf{x} = \xi^2$ and $\mathbf{y} = \xi$, for $-\infty < \xi < \infty$. (ii) Write the ode for the characteristic curves, in terms of some parameter (say, s) along each curve. (iii) Solve the ode for the characteristics, with the condition that $\mathbf{x} = \xi^2$ and $\mathbf{y} = \xi$, for $s = 0$.
- B.** Draw the characteristics, in the x - y plane, that you just computed. Which region of the plane do the curves cover? What happens with the characteristic corresponding to $\xi = 0$?
- C.** Solve the ode that ψ satisfies along each characteristic. Eliminate the parameters ξ and s in terms of x and y , and write an explicit formula for the solution $\psi = \psi(x, y)$ to (2.11).
- D.** Where is the solution ψ defined? **Hint.** *Be careful with your answer here! What happens with ψ along each characteristic, far enough from Γ ? What happens with the $\xi = 0$ characteristic?*

THE END.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.311 Principles of Applied Mathematics
Spring 2014

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.