

Course 18.312: Algebraic Combinatorics

HW # 6

Will be collected March 20, 2009

Due to the length of MT # 1, I am grading the midterm as if it were out of 55 points, and assigning problems #3 and #4 in lieu of homework this week. This problem set will either count towards your homework grade or your Midterm #1 score in the final grade calculation. It may be worth up to 45 midterm points, with the possibility of 10 bonus points. You may collaborate, but please list your collaborators as usual. You may not obtain solutions from other sources including computer software.

- 3) Define a family of graphs, we call them pinwheel graphs, as follows: Let PW_T be the graph on $2T + 1$ vertices

$$u_0 \cup \{v_1, \dots, v_T\} \cup \{w_1, \dots, w_T\}$$

such that there is an edge between u_0 and every other vertex and v_i is connected to w_j if and only if $i = j$.

(10 points) a) Compute the eigenvalues of the adjacency matrix $A(PW_T)$ in terms of T .

Hint: One way to approach this problem is to use symmetry to find a large set of linearly independent eigenvectors with first entry zero, and use combinatorial formulas to deduce the remaining eigenvalues.

(5 points) b) Describe the set of T such that PW_T is an integral graph.

- 4) Define the following family of posets: For all integers $n \geq 1$, P_n is the poset, ordered by inclusion, consisting of subsets $\{i_1, i_2, \dots, i_{2k}\} \subset \{1, 2, 3, \dots, 2n\}$ satisfying

$$0 < i_1 < i_2 < \dots < i_{2k} < 2n + 1$$

and

$$i_1, (i_2 - i_1), \dots, (i_{2k} - i_{2k-1}), (2n + 1) - i_{2k} \quad \text{are all odd.}$$

Notice that all elements are subsets with an even number of elements and that the rank of an element S is $|S|/2$. We let $\hat{0}$ denote \emptyset , the unique element of rank 0 and $\hat{1}$ denote $\{1, 2, \dots, 2n\}$, the unique element of rank n .

P_n also has the property, which you do not need to show, that if $\text{rank}(S) = k$, then the interval $[\hat{0}, S]$ is isomorphic to P_k .

a) Draw the Hasse Diagram for P_3 .

(5 points) b) Compute the Möbius function $\mu(\hat{0}, \hat{1})$ for P_3 .

(5 points) c) Compute the total number of elements in P_n .

(5 points) d) Show that the number of elements of rank k in P_n is $\binom{n+k}{2k}$.

It is easy to show, you do not need to, that if S is of rank k , then the cardinality $\#\{S' : S \text{ covers } S'\}$ is a function $f(k)$ only depending on k . In other words, it is the same number, regardless of the choice of S or choice of P_n .

(5 points) e) What is $f(k)$?

(5 points) f) Using the above, deduce a formula for the number of maximal chains in P_n and prove it.

(5 points) g) Using the above or otherwise, compute $\mu(\hat{0}, \hat{1})$ for P_4 .

(Bonus 10 points) Deduce a formula for $\mu(\hat{0}, \hat{1})$ in P_n and prove it.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.312 Algebraic Combinatorics
Spring 2009

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.