

18.445 Introduction to Stochastic Processes

Lecture 20: Poisson process

Hao Wu

MIT

29 April 2015

Random point process

A random point process is a countable random set of points of the real line which corresponds to the sequence of the times of occurrence of some event. For instance, the arrival times of customers.

Definition

A random point process on \mathbb{R}_+ is a sequence of random variables $(T_n)_{n \geq 0}$ such that

- $0 = T_0 < T_1 < T_2 < \dots$
- $\lim_n T_n = \infty$

Definition

The interevent sequence : $S_n = T_n - T_{n-1}$ for $n \geq 1$.

The counting process : For $(a, b] \subset \mathbb{R}_+$, define

$$N(a, b] = \sum_{n \geq 1} 1_{(a, b]}(T_n)$$

Counting process

Definition

The counting process : For $(a, b] \subset \mathbb{R}_+$, define

$$N(a, b] = \sum_{n \geq 1} 1_{(a, b]}(T_n).$$

In particular, set $N_t = N(0, t]$. Then

- $N_0 = 0$
- $N(a, b] = N_b - N_a$
- $t \mapsto N_t$ is right-continuous

Poisson process

Definition

A point process N on \mathbb{R}_+ is called a Poisson process with intensity $\lambda > 0$ if

- For any $k \geq 1$, any $0 \leq t_1 \leq t_2 \leq \dots \leq t_k$, the random variables $N(t_i, t_{i+1}]$, $i = 1, \dots, k - 1$ are independent.
- For any interval $(a, b] \subset \mathbb{R}_+$, the variable $N(a, b]$ is a Poisson random variable with mean $\lambda(b - a)$, i.e.

$$\mathbb{P}[N(a, b] = k] = e^{-\lambda(b-a)} \frac{(\lambda(b-a))^k}{k!}.$$

Theorem

The interevent sequence $(S_n)_{n \geq 1}$ of a Poisson process with intensity λ is i.i.d. with exponential distribution of parameter λ .

Poisson process — Markov property

Theorem (Markov property)

Let $(N_t)_{t \geq 0}$ be a Poisson process. Then, $\forall s \geq 0$,

- the process $(N_{t+s} - N_s)_{t \geq 0}$ is also a Poisson process
- and it is independent of $(N_u)_{u \leq s}$

Theorem (Strong Markov property)

Let $(N_t)_{t \geq 0}$ be a Poisson process. Suppose that T is a stopping time, then conditional on $[T < \infty]$,

- the process $(N_{t+T} - N_T)_{t \geq 0}$ is also a Poisson process
- and it is independent of $(N_u)_{u \leq T}$

Poisson process — Superposition

Theorem

Let $(N^i)_{i \geq 1}$ be a family of independent Poisson processes with respective positive intensities $(\lambda_i)_{i \geq 1}$. Then

- two distinct Poisson processes in this family have no points in common
- if $\sum_{i \geq 1} \lambda_i = \lambda < \infty$, then $N_t = \sum_{i \geq 1} N_t^i$ defines the counting process of a Poisson process with intensity λ .

Theorem

In this situation of the above theorem with $\sum \lambda_i = \lambda < \infty$. Denote by Z the first event time of $N = \sum N^i$ and by J the index of the Poisson process responsible for it. Then

$$\mathbb{P}[J = i, Z \geq a] = \mathbb{P}[J = i] \times \mathbb{P}[Z \geq a] = \frac{\lambda_i}{\lambda} e^{-\lambda a}.$$

Poisson process — Characterization

Theorem

Let $(X_t)_{t \geq 0}$ be an increasing right-continuous process taking values in $\{0, 1, 2, \dots\}$ with $X_0 = 0$. Let $\lambda > 0$. Then the following statements are equivalent.

- $(X_t)_{t \geq 0}$ is a Poisson process with intensity λ .
- X has independent increments, and as $\epsilon \downarrow 0$, uniformly in t , we have

$$\mathbb{P}[X_{t+\epsilon} - X_t = 0] = 1 - \lambda\epsilon + o(\epsilon);$$

$$\mathbb{P}[X_{t+\epsilon} - X_t = 1] = \lambda\epsilon + o(\epsilon).$$

- X has independent and stationary increments, and for all $t \geq 0$ we have $X_t \sim \text{Poisson}(\lambda t)$.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.445 Introduction to Stochastic Processes

Spring 2015

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.