

MIT OpenCourseWare
<http://ocw.mit.edu>

18.969 Topics in Geometry: Mirror Symmetry
Spring 2009

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

MIRROR SYMMETRY: LECTURE 5

DENIS AUROUX

1. GROMOV-WITTEN INVARIANTS

Recall that if (X, ω) is a symplectic manifold, J an almost-complex structure, $\beta \in H_2(X, \mathbb{Z})$, $\overline{\mathcal{M}}_{g,k}(X, J, \beta)$ is the set of (possibly nodal) J -holomorphic maps to X of genus g representing class β with k marked points up to equivalence. This is not a nice moduli space, but does have a fundamental class $[\overline{\mathcal{M}}_{g,k}(X, J, \beta)] \in H_{2d}(\overline{\mathcal{M}}_{g,k}(X, J, \beta), \mathbb{Q})$, where $2d = \langle c_1(TX), \beta \rangle + 2(n-3)(1-g) + 2k$. We further have an evaluation map $\text{ev} = (\text{ev}_1, \dots, \text{ev}_n) : \overline{\mathcal{M}}_{g,k}(X, J, \beta) \rightarrow X^k$, $(\Sigma, z_1, \dots, z_k, u) \mapsto (u(z_1), \dots, u(z_k))$. Then the Gromov-Witten invariants are defined for $\alpha_1, \dots, \alpha_k \in H^*(X)$, $\sum \deg \alpha_i = 2d$ by

$$(1) \quad \langle \alpha_1, \dots, \alpha_k \rangle_{g,\beta} = \int_{[\overline{\mathcal{M}}_{g,k}(X, J, \beta)]} \text{ev}_1^* \alpha_1 \wedge \dots \wedge \text{ev}_k^* \alpha_k \in \mathbb{Q}$$

Or dually, for $\alpha_i = PD(C_i)$, $\#(\text{ev}_*[\overline{\mathcal{M}}_{g,k}(X, J, \beta)] \cap (C_1 \times \dots \times C_k)) \in \mathbb{Q}$.

For a Calabi-Yau 3-fold, we're interested in $g = 0, k = 3$, so $\Sigma = (S^2, \{0, 1, \infty\})$. For $\deg \alpha_i = 2, \alpha_i = PD(C_i)$, C_i cycles transverse to the evaluation map, we have

$$(2) \quad \langle \alpha_1, \alpha_2, \alpha_3 \rangle_{0,\beta} = \#\{u : S^2 \rightarrow X \text{ } J\text{-hol. of class } \beta, \\ u(0) \in C_1, u(1) \in C_2, u(\infty) \in C_3\} / \sim$$

Reparameterization acts transitively on triples of points, so

$$(3) \quad \langle \alpha_1, \alpha_2, \alpha_3 \rangle_{0,\beta} = (C_1 \cdot \beta)(C_2 \cdot \beta)(C_3 \cdot \beta) \#\{u : S^2 \rightarrow X \text{ } J\text{-hol. of class } \beta\} / \sim \\ = \left(\int_{\beta} \alpha_1 \right) \left(\int_{\beta} \alpha_2 \right) \left(\int_{\beta} \alpha_3 \right) \cdot \#[\overline{\mathcal{M}}_{0,0}(X, J, \beta)]$$

We denote by $N_{\beta} \in \mathbb{Q}$ the latter number $\#[\overline{\mathcal{M}}_{0,0}(X, J, \beta)]$. This works when $\beta \neq 0$: when $\beta = 0$, we instead obtain

$$(4) \quad \langle \alpha_1, \alpha_2, \alpha_3 \rangle_{0,0} = \int_X \alpha_1 \wedge \alpha_2 \wedge \alpha_3$$

1.1. **Yukawa coupling.** Physicists write this as

$$(5) \quad \langle \alpha_1, \alpha_2, \alpha_3 \rangle = \int_X \alpha_1 \wedge \alpha_2 \wedge \alpha_3 + \sum_{0 \neq \beta \in H_2(X, \mathbb{Z})} \langle \alpha_1, \alpha_2, \alpha_3 \rangle_{0, \beta} e^{2\pi i \int_\beta B + i\omega}$$

We want to ignore issues of convergence, and so treat this as a formal power series

$$(6) \quad \langle \alpha_1, \alpha_2, \alpha_3 \rangle = \int_X \alpha_1 \wedge \alpha_2 \wedge \alpha_3 + \sum_{\beta \neq 0} \langle \alpha_1, \alpha_2, \alpha_3 \rangle_{0, \beta} q^\beta \in \Lambda$$

where Λ is the completion of the group ring $\mathbb{Q}[H_2(X, \mathbb{Z})] = \{\sum a_i q^{\beta_i} \mid a_i \in \mathbb{Q}, \beta_i \in H_2\}$. Specifically, we allow infinite sums provided that $\int_{\beta_i} \omega \rightarrow +\infty$.

1.2. **Quantum cohomology.** This is new product structure on $H^*(X)$ deformed by this coupling. Namely, pick a basis (η_i) of $H^*(X)$, (η^i) the dual basis, i.e. $\int_X \eta_i \wedge \eta^j = \delta_{ij}$. Set

$$(7) \quad a_1 * a_2 = \sum_i \langle \alpha_1, \alpha_2, \eta^i \rangle \eta_i = \alpha_1 \wedge \alpha_2 + \sum_{\beta \neq 0} \langle \alpha_1, \alpha_2, \eta^i \rangle_{0, \beta} q^\beta \eta_i$$

Definition 1. *The quantum cohomology of X is $QH^*(X) = (H^*(X; \Lambda), *)$.*

Theorem 1. *This is an associative algebra.*

The proof of this relies on understanding the relationship between 4 point GW invariants and various 3 point ones.

1.3. **Kähler moduli.** We can view q as the coordinates on a Kähler moduli space: for (X, J) -complex, the Kähler cone $\mathcal{K}(X, J) = \{[\omega] \mid \omega \text{ Kähler}\} \subset H^{1,1}(X) \cap H^2(X, \mathbb{R})$ is a open, convex cone. Its real dimension is $h^{1,1}(X)$, and we can make it a complex manifold by adding the ‘‘B-field’’.

Definition 2. *Let (X, J) be a Calabi-Yau 3-fold with $h^{1,0} = 0$ (so $h^{2,0} = 0$ and $H^{1,1} = H^2$). Then the complexified Kähler moduli space is*

$$(8) \quad \begin{aligned} \mathcal{M}_{Kah} &= (H^2(X, \mathbb{R}) + i\mathcal{K}(X, J)) / H^2(X, \mathbb{Z}) \\ &= \{[B + i\omega], \omega \text{ Kahler}\} / H^2(X, \mathbb{Z}) \end{aligned}$$

Choose a basis (e_i) of $H^2(X, \mathbb{Z})$, $e_1, \dots, e_m \in \overline{\mathcal{K}(X, J)}$ (which exists by openness). We can write $[B + i\omega] = \sum t_i e_i$, $t_i \in \mathbb{C}/\mathbb{Z}$, so we have coordinates on \mathcal{M}_{Kah} given by $q_i = \exp(2\pi i t_i)$. Thus, \mathcal{M}_{Kah} is an open subset of $(\mathbb{C}^*)^m$ which contains $(\mathbb{D}^*)^m$, where $\mathbb{D}^* = \{q \mid 0 < |q| < 1\}$.

We now can associate q^β to $q_1^{d_1} \cdots q_m^{d_m}$, where $d_i = \int_\beta e_i$ for $e_i \geq 0$ integers (it is an integer cohomology class integrated against an integer homology class): explicitly, $q_1^{d_1} \cdots q_m^{d_m} = \exp(2\pi i \sum d_i t_i) = \exp(2\pi i \int_\beta B + i\omega)$. We can view $\langle \alpha_1, \alpha_2, \alpha_3 \rangle$ as a power series in the q_i , though we still do not know about convergence.

1.4. **Gromov-Witten invariants vs. numbers of curves.** We have, for $\alpha_1, \alpha_2, \alpha_3 \in H^2(X)$,

$$(9) \quad \begin{aligned} \langle \alpha_1, \alpha_2, \alpha_3 \rangle &= \int_X \alpha_1 \wedge \alpha_2 \wedge \alpha_3 + \sum_{\beta \neq 0} \langle \alpha_1, \alpha_2, \alpha_3 \rangle_{0, \beta} q^\beta \\ &= \int_X \alpha_1 \wedge \alpha_2 \wedge \alpha_3 + \sum_{\beta \neq 0} \left(\int_\beta \alpha_1 \right) \left(\int_\beta \alpha_2 \right) \left(\int_\beta \alpha_3 \right) N_\beta q^\beta \end{aligned}$$

This is much like our formula from the first class, except the latter term had the form $n_\beta \frac{q^\beta}{1-q^\beta}$ and n_β as the number of “rational curves of class β ”. The discrepancy comes from the existence of multiple covers. Let $C \subset X$ be an embedded rational curve in a Calabi-Yau 3-fold. A theorem of Grothendieck says that a holomorphic bundle over \mathbb{P}^1 splits as $\bigoplus \mathcal{O}_{\mathbb{P}^1}(d_i)$, where $\mathcal{O}(d)$ is the sheaf whose sections are homogeneous degree d holomorphic functions on \mathbb{C}^2 and $\mathcal{O}(-1)$ is the tautological bundle. Writing $NC \cong \mathcal{O}_{\mathbb{P}^1}(d_1) \oplus \mathcal{O}_{\mathbb{P}^1}(d_2)$, we obtain

$$(10) \quad 0 = c_1(TX)[C] = c_1(NC)[C] + c_1(TC)[C] = d_1 + d_2 + 2$$

so $d_1 + d_2 = -2$. The “generic case” is $d_1 = d_2 = -1$, in which case C is automatically regular as a J -holomorphic curve. The contribution of C to the Gromov-Witten invariant $N_{[C]}$ is precisely 1. On the other hand, there is a component $\mathcal{M}(kC) \subset \mathcal{M}_{0,0}(X, J, k[C])$ consisting of k -fold covers of C . What is $\#\mathcal{M}(kC)$?

Theorem 2. *If $NC \cong \mathcal{O}(-1) \oplus \mathcal{O}(-1)$, then the contribution of C to $N_{k[C]}$ is $\frac{1}{k^3}$.*

There are various proofs, all of which are somewhat difficult. For instance, Voisin shows that \exists perturbed $\bar{\partial}$ -equations $\bar{\partial}_J u = \nu(z, u(z))$ s.t. the moduli space $\tilde{M}_3(kC)$ (of perturbed J -holomorphic maps with 3 marked points representing $k[C]$ and whose image lies in a neighborhood of C) is smooth and has real dimension 6. Moreover, $(\text{ev}_1 \times \text{ev}_2 \times \text{ev}_3)_*[\tilde{M}_3(kC)] = [C \times C \times C] \in H_6(X \times X \times X)$. Then the contribution of C to $\langle \alpha_1, \alpha_2, \alpha_3 \rangle_{0, k[C]}$ is

$$(11) \quad \int_{\text{ev}_*[\tilde{M}_3]} \alpha_1 \times \alpha_2 \times \alpha_3 = \left(\int_C \alpha_1 \right) \left(\int_C \alpha_2 \right) \left(\int_C \alpha_3 \right) = \frac{1}{k^3} \left(\int_{kC} \alpha_1 \right) \left(\int_{kC} \alpha_2 \right) \left(\int_{kC} \alpha_3 \right)$$

We expect that (*) $N_\beta = \sum_{\beta=k\gamma} \frac{1}{k^3} n_\gamma$.

Remark. We do not know if n_γ is what we think it is, but we use this formula as a definition; see the Gopakumar-Vafa conjecture, which claims that $n_\gamma \in \mathbb{Z}$, and the theory of Donaldson-Thomas invariants and MNOP conjectures.

Assuming (*), we have

$$\begin{aligned}
 (12) \quad \sum_{\beta} \left(\int_{\beta} \alpha_1 \right) \left(\int_{\beta} \alpha_2 \right) \left(\int_{\beta} \alpha_3 \right) N_{\beta} q^{\beta} &= \sum_{k, \gamma} \left(\int_{k\gamma} \alpha_1 \right) \left(\int_{k\gamma} \alpha_2 \right) \left(\int_{k\gamma} \alpha_3 \right) \frac{n_{\gamma}}{k^3} q^{k\gamma} \\
 &= \sum_{\gamma} \left(\int_{\gamma} \alpha_1 \right) \left(\int_{\gamma} \alpha_2 \right) \left(\int_{\gamma} \alpha_3 \right) n_{\gamma} \sum_{k \geq 1} k^{k\gamma}
 \end{aligned}$$

Where we are headed: we correspond this pairing to

$$(13) \quad \langle \theta_1, \theta_2, \theta_3 \rangle = \int_X \Omega \wedge (\nabla_{\theta_1} \nabla_{\theta_2} \nabla_{\theta_3} \Omega)$$

on $H^{2,1}(\check{X})$.